

A New Host Record *Utetheisa pulchella* (Linnaeus, 1758) (Lepidoptera: Erebidæ) for *Exorista xanthaspis* (Wiedemann, 1830) (Diptera: Tachinidae) from Turkey

Halil BOLU¹

Turgut ATAY^{2*}

Kenan KARA³

Hakan ÇELİK⁴

^{1,4}Dicle University, Faculty of Agriculture, Department of Plant Protection, Diyarbakır, TURKEY

^{2,3}Tokat Gaziosmanpasa University, Faculty of Agriculture, Department of Plant Protection, Tokat, TURKEY

e-mails: ¹besni@dicle.edu.tr, ^{2*}turgut.atay@gop.edu.tr, ³kenan.kara@gop.edu.tr,

⁴144061018a@gmail.com

ORCID IDs: ¹0000-0001-5488-0056, ²0000-0002-9074-0816, ³0000-0003-0439-5639

⁴0000-0002-2318-3474

ABSTRACT

Exorista xanthaspis (Wiedemann, 1830) (Diptera: Tachinidae) specimens are reared from the larvae of *Utetheisa pulchella* (Linnaeus, 1758) (Lepidoptera: Erebidæ) collected in Batman province. *U. pulchella* is recorded for the first time as host of this parasitoid. Some additional information about the reared species and its host is also provided.

Key words: *Exorista xanthaspis*, new host record, *Utetheisa pulchella*, Turkey.

INTRODUCTION

Tachinid flies (Diptera: Tachinidae) are important in terms of biological control because their larvae develop as parasitoids in insects and other arthropods. The majority of hosts are caterpillars of Lepidoptera. Other hosts belong to the orders Coleoptera, Hemiptera, Hymenoptera, Orthoptera and Diptera (Grenier, 1988; Stireman, O'Hara, & Wood, 2006; Tschorsnig, 2017). Many hosts are still unknown. Recently, the most comprehensive host-parasitoid catalogues about Turkey and the Palaearctic region were prepared by Kara & Tschorsnig (2003) and Tschorsnig (2017), respectively.

The tachinid species *Exorista xanthaspis* (Wiedemann, 1830) has a broad host range in the Palaearctic region. Lasiocampidae, Lymantriidae and Noctuidae (Lepidoptera) are the usual host families of this tachinid. Other lepidopterous host families in the same region are Arctiidae, Epicopeiidae, Pieridae, Pyralidae, Sphingidae and Thaumetopoeidae (Tschorsnig, 2017). Also, Noctuidae is a common host family in the Afrotropical and Oriental regions. Other host families of this tachinid in the Oriental region are Arctiidae and Hyblaeidae (Lepidoptera) (Crosskey, 1976; 1984).

There are only few records on other tachinid parasitoids of *U. pulchella*. These are *Exorista segregata* (Rondani, 1859) (Kugler, 1980) and *Tachina praeceps* Meigen, 1824 (Herting, 1960). There were no published records of Tachinidae reared from *U. pulchella* in Turkey.

MATERIAL AND METHODS

Thirty-three larvae of the *Utetheisa pulchella* (Lep.: Erebiidae) were collected on *Heliotropium ellipticum* (Boraginaceae) in Batman and Diyarbakır provinces in 2018. They were brought to the laboratory with their food-plants for rearing and transferred to separate cages and checked daily.

Male terminalia of the reared parasitoids were prepared following the method described by O'Hara (2002). The dissected terminalia were examined with a Leica M205 C stereoscopic microscope and are preserved in small plastic vials with glycerol. Images were taken using a Leica MC 170 digital camera mounted on a Leica M205 C stereoscopic microscope, and processed with Helicon Focus Pro software. The keys of Herting (1975) and Tschorsnig & Herting (1994) were used for the identification of the species. The nomenclature of the tachinids follows Herting & Dely-Draskovits (1993). The lepidopterous host was identified by Felipe Gil-T (Granada, Spain). The specimens are deposited at the Plant Protection Museum of the Tokat Gaziosmanpaşa University, Agricultural Faculty, Plant Protection Department, Tokat, Turkey.

RESULTS

Identity, distribution, and some additional information of tachinid and host are as follows:

***Utetheisa pulchella* (Linnaeus, 1758) (Lepidoptera: Erebidæ)**

The Crimson-speckled moth *U. pulchella*, which attacks some cultivated plants, is a polyphagous leaf feeder pest (Mekhlif, 2012).

Distribution: Europe (Olafsson, et al. 2019); India (Dubatolov, 2010; Bhatt, 2016; Biswas, Modak, Mazumder, & Mitra, 2016), Libya (El-Maghrabi & Amin, 2007) Iraq (Mekhlif, 2012), Turkey: Çukurova Deltası (Aydın, 2006), Şanlıurfa (Beyarslan, Gözüaık, & Özgen, 2014), Şanlıurfa (Kemal & Koak, 2017).

Host plants: *Crotalaria juncea* L. (Beyarslan et al, 2014), *Crotalaria burhia* Buch.-Ham. (Fabaceae) (Pandey, Pande, & Kaul, 1971); *Heliotrobium ramosissimum* (Lehm.) (Boraginaceae), *Launaea cassiniana* (Jaub. & Spach) (Asteraceae), *Gossypium* sp. (Malvaceae), *Ricinus communis* L. (Euphorbaceae), *Lawsonia incamis* (Lythraceae), *Medicago sativa* L. (Fabaceae), *Lycopersicum esculentum* Mill., *Solanum melongena* L., *Withania somnifera* (L.) (Solanaceae) (AL-Ahmadi & Salem, 1995), *Myosotis* sp. (Boraginaceae) (Becker & Scott 2002), *Heliotropium ovalifolium* Forssk. (Boraginaceae) (Bhatt, 2016).

Material examined: Collected in Batman: Hasankeyf, 2.09.2018, N 37°42.43.92', E 41°24.38.14', 516 m, on *Heliotropium ellipticum* Ledeb. (Boraginaceae); in Diyarbakır: Sur, 7.09.2018, N 37°55.32.62', E 40°15.32.62', 613 m, on *H. ellipticum* (Fig. 1).



Fig. 1. Larvae of *Utetheisa pulchella*.

***Exorista xanthaspis* (Wiedemann, 1830) (Tachinidae: Exoristinae)**

Distribution: Caucasus, East Siberia, Mongolia, Soviet Middle Asia, Sudan (Herting & Dely-Draskovits, 1993), Israel, India, Indonesia, Madagascar, Taiwan, Yemen (O'Hara & Cerretti, 2016), East, South and West Europe (Tschorsnig, et al. 2004), Turkey: Erzurum (Doğanlar, 1975), Diyarbakır (Efil & Kara, 2004), Mardin (Gözüaık & Mart, 2009), Southeastern Anatolia Region (Gözüaık, Mart, & Kara, 2009).

Hosts in Turkey: *Aporia crataegi* (Lep.: Pieridae) (Kansu, 1955), *Simyra dentinosa* (Lep.: Noctuidae) (Doğanlar, 1982), *Spodoptera exigua* (Lep.: Noctuidae) (Steiner 1937; Efil & Kara 2004; Gözüaık et al, 2009; Gözüaık & Mart 2009). *Utetheisa pulchella* is a new host species for this tachinid in the world.

Reared specimens (date of adult emergence): 1♀ (18.09.2018); 1♂, 1♀ (14.10.2018).

Although the specimens of *E. xanthaspis* were reared from *U. pulchella* collected from Batman (Hasankeyf), they could not be reared from those collected from Diyarbakır (Sur).

Differential diagnosis: *Exorista xanthaspis* shows many external morphological characters similar to *E. civilis* (Rondani, 1859). For the safe distinction of the two species a study of the male terminalia is recommended (Herting, 1975).

- Syncercus with a large basal part and a short, blunt, slightly flattened top. Surstylus almost in line with the syncercus. Aedeagus nearly continuously bent (Fig. 2)*E. civilis*

- Syncercus with a smaller basal part and a long, laterally compressed tip. Surstylus semi-erect. Aedeagus dorsally with a hump at about mid-length (Fig. 3)*E. xanthaspis*




Fig. 2: Male terminalia of *Exorista civilis* a. Syncercus b. Surstyli c. Aedeagus (Herting, 1975).



Fig. 3: Male terminalia of *Exorista xanthaspis* a. Syncercus b. Surstyli c. Aedeagus.

ACKNOWLEDGEMENTS

We are grateful to Dr. H.-P. Tschorsnig (Staatliches Museum für Naturkunde,

A New Host Record Utetheisa pulchella for Exorista xanthaspis

Stuttgart, Germany) for providing several literature and information on host-parasitoid couple and to Felipe Gil-T (Granada, Spain) for identification of *Utetheisa pulchella*.

REFERENCES

- AL-Ahmadi, A.Z. & Salem, M.M. (1995). *Entomofauna of Saudia Arabia. Part II Checklist of Phytophagous Insects*. Academic Publication and Press, King Saud University Press. Riyadh, Saudi Arabia.
- Aydın, G. (2006). Çukurova Deltası'nda Böceklerin Sürdürülebilir Alan Kullanımında Biyolojik Gösterge Olarak Değerlendirilme Olanakları. Doktora Tezi, Çukurova Üniversitesi Fen Bilimleri Enstitüsü, Bitki Koruma Anabilim Dalı, Adana, 307.
- Becker, V.O. & Scott, E.M. (2002). The large moths of Guana Island, British Virgin Islands: A survey of efficient colonizers (Sphingidae, Notodontidae, Noctuidae, Arctiidae, Geometridae, Hyblaeidae, Cossidae). *Journal of the Lepidopterists Society*, 56(1), 9-44.
- Beyarslan, A., Gözüaçık, C., & Özgen, İ. (2014). First research on Braconinae fauna of South-eastern Anatolia region with new localities of Turkey (Hymenoptera: Braconidae). *Entomofauna Zeitschrift für Entomologie*, 35(10), 177-204.
- Bhatt, N.A. (2016). *Heliotropium ovalifolium* Forsk., a weed, as a host of *Utetheisa pulchella* L. (Lepidoptera: Erebiidae). *Biotic Environment, formerly Insect Environment*, 21(4), 61-63.
- Biswas, O., Modak, B.K., Mazumder, A., & Mitra, B. (2016). Moth (Lepidoptera: Heterocera) diversity of Sunderban biosphere reserve, India and their pest status to economically important plants. *Journal of Entomology and Zoology Studies*, 4(2), 13-19.
- Crosskey, R.W. (1976). A taxonomic conspectus of the Tachinidae (Diptera) of the Oriental Region. *Bulletin of the British Museum (Natural History). Entomology Supplement*, 26, 1-357.
- Crosskey, R.W. (1984). Annotated keys to the genera of Tachinidae (Diptera) found in tropical and southern Africa. *Annals of the Natal Museum*, 26(1), 189-337.
- Doğanlar, M. (1975). *Erzurum bölgesinde önemli lepidopter tırtıllarında bulunan Tachinidae sinekleri ve bunların kısa biyolojileri*. Atatürk Üniversitesi. Erzurum, 136pp.
- Doğanlar, M. (1982). Some parasitic flies from Eastern Anatolia I. (Diptera: Tachinidae, Exoristinae). *Türkiye Bitki Koruma Dergisi*, 6(3), 75-79.
- Dubatolov, V.V. (2010). Tiger moths of Eurasia (Lepidoptera: Arctiinae). *Neue Entomologische Nachrichten*, 65, 1-106.
- Efil, L. & Kara, K. (2004). Tachinid parasitoids (Diptera: Tachinidae) of *Spodoptera exigua* in cotton fields in Diyarbakir, Turkey. *Phytoparasitica*, 32(4), 363-366.
- El-Maghrabi, M. S. & Amin, A. H. (2007). List of the Lepidoptera insects surveyed in El-Beida Area, with their world distribution, host plants and notes on taxonomy. *Journal of Science and its Applications*, 1(2), 21-31.
- Gözüaçık, C. & Mart, C. (2009). Determination of natural parasitization rates of some pests of Lepidoptera larvae in corn (*Zea mays* L.) in the Southeastern Anatolia Region. *Bitki Koruma Bülteni*, 49 (3), 107-116.
- Gözüaçık, C., Mart, C. & Kara, K. (2009). Parasitoids of several lepidopterous pests in maize plantations in the Southeast Anatolian Region of Turkey. *Turkish Journal of Zoology*, 33, 475-477.
- Grenier, S. (1988). Applied biological control with tachinid flies (Diptera, Tachinidae). A review, *Anzeiger für schädlingkunde, Pflanzenschutz, Umweltschutz*, 61(3), 49-56.
- Herting, B. (1960). *Biologie der westpaläarktischen Raupenfliegen (Dipt., Tachinidae)*. Monographien zur angewandten Entomologie, Hamburg und Berlin, 188.
- Herting, B. (1975). Nachträge und Korrekturen zu den von Meigen und Rondani beschriebenen Raupenfliegen (Dipt. Tachinidae). *Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie)*, 271, 1-13.
- Herting, B. & Dely-Draskovits, Á. (1993). Family Tachinidae. In A., Soós, L. Papp, (Eds.). *Catalogue of palaearctic Diptera. Anthomyiidae-Tachinidae* Budapest, 118-458.

- Kansu, A. (1955). *Orta Anadolu meyve ağaçlarına zarar veren bazı Makrolepidoptera türlerinin evsafı ve kısa biyolojileri hakkında araştırmalar*. Ziraat Vekaleti Neşriyat ve Haberleşme Müdürlüğü, Ankara, 204
- Kara, K. & Tschorsnig, H.P. (2003). Host Catalogue for the Turkish Tachinidae (Diptera). *Journal of Applied Entomology*, 127, 465-476.
- Kemal, M. & Koçak, A.Ö. (2017). On the vernal Lepidoptera fauna of Nizip-Birecik districts-Euphrates region in South Turkey. *Cesa News Centre for Entomological Studies* Ankara, 22.
- Kugler, J. (1980). New taxa of Tachinidae (Diptera) with a list of the species from Israel and Adjacent Territories. *Israel Journal of Entomology*, 13, 27-60.
- Mekhlif, A.F. (2012). Extract bioefficiency of five *Euphorbia* spp. (Euphorbaceae) on crimson-speckled moth, *Utetheisa pulchella* L. (Lepidoptera, Acritidae) growth and development. *Rafidain Journal of Science*, 23(4), 23-32.
- O'Hara, J.E. (2002). Revision of the Polideini (Tachinidae) of America North of Mexico. *Studia Dipterologica*, Supplement, 10, 170pp.
- O'Hara, J.E. & Cerretti, P. (2016). Annotated catalogue of the Tachinidae (Insecta, Diptera) of the Afrotropical region, with the description of seven new genera. *ZooKeys*, 575, 1-344.
- Olafsson, E., Kullberg, J., Jensen, J.K., Ronkay, L., Corley, M., Sammut, P.M., Prins, W.D., Tokar, Z., & Karsholt, E.J. (2019). Erebidae. Fauna Europaea Service, <http://www.faunaeur.org>. (16.01.2019).
- Pandey, S.N., Pande, Y.D., & Kaul, C.L. (1971). An alternate host plant of *Utetheisa pulchella* L. (Arctiidae: Lepidoptera). *Indian Journal of Entomology*, 32(3), 277.
- Steiner, P. (1937). Beiträge zur Kenntnis der Schädlingfauna Kleinasien III. *Laphygma exigua* Hb., ein Groß-Schädling der Zuckerrübe in Anatolien. *Zeitschrift für Angewandte Entomologie*, 23, 177-222.
- Stireman, J.O., O' Hara, J.E., & Wood, D.M. (2006). Tachinidae: Evolution, behavior and ecology. *Annual Review of Entomology*, 51(1), 525-555.
- Tschorsnig, H.P. (2017). Preliminary host catalogue of Palaearctic Tachinidae (Diptera). 480pp www.nadsdiptera.org/Tach/WorldTachs/CatPalHosts/Home.html. (09.11.2018).
- Tschorsnig H.P. & Herting, B. (1994). *Die Raupenfliegen (Diptera: Tachinidae) Mitteleuropas: Bestimmungstabellen und Angaben zur Verbreitung und Ökologie der einzelnen Arten*. Stuttgarter Beiträge zur Naturkunde (A) 506: 1-170. Online Authorized Version of English Translation by Rayner R. & Raper C.: Tschorsnig H.P. and Herting B., 2001: The Tachinids (Diptera:Tachinidae) of Central Europe: Identification Keys for the Species and Data on Distribution and Ecology, <http://tachinidae.org.uk/site/downloads.php>. 12.06.2006.
- Tschorsnig, H.P., Richter, V.A., Cerretti, P., Zeegers, T., Bergström, C., Vaňhara, J., Van de Weyer, G., Bystrowski, C., Raper, C., Ziegler, J. & Hubenov, Z. (2004). Tachinidae. Fauna Europaea Service, <http://www.faunaeur.org>. (29.12.2018).

Received: January 03, 2019

Accepted: September 16, 2019